

Delirium

A Brief Introduction for the Braidwood Inquiry



**SHAOHUA LU MD FRCPC
ADDICTION PSYCHIATRIST
CLINICAL ASSISTANT PROFESSOR UBC**

Delirium Definition and Presentations



- **Delirium is also known as**
 - Acute confusional state
 - Acute organic brain syndrome
 - Acute mental agitation
 - Acute brain syndrome
 - Acute brain failure
 - Toxic encephalopathy

Delirium Definition and Presentations



- **DSM-IV-TR Criteria**

- Disturbance of the consciousness with reduced ability to focus or sustain attention
- A change in cognition or the development of a perceptual disturbance that is not better accounted for by a pre-existing dementia
- The disturbance develops over a short time and tends to fluctuate during the course of the day

Delirium Definition and Presentations



- **Generalized Cognitive and Brain Dysfunction**
- **Impaired Level of Consciousness**
 - Drowsiness, agitation, disrupted sleep-wake cycle
- **Attentional Deficits**
 - Disorientation
 - Poor attention and concentration
- **Language Deficits**
 - Impaired comprehension and speech
- **Memory Deficits**
 - Impaired short term and working memory

Delirium Definition and Presentations



- **Executive Cognitive Function Impairments**
 - Decreased ability to plan, organize and execute tasks
 - General disorganization
 - Impaired insight
- **Thought Disturbances**
 - Delusional thinking or illogical thought process
 - Potentials for bizarre behaviours
- **Disturbed Affects**
 - Labile emotional expression
 - Anxiety
 - Irritability

Delirium Definition and Presentations



- **Perceptual Disturbances**
 - Misinterpretation of external and internal stimuli
 - Hallucination (visual, auditory or tactile)
- **Psychomotor Changes**
 - Agitation and restlessness
 - Defensive aggression
 - (Excited delirium: extreme agitation during delirium)
- **Autonomic Instability**
 - Sweating, rapid breathing, increased heart rate and unstable blood pressure

Delirium Definition and Presentations



- Often acute onset
- Fluctuating course
- A medical syndrome
- Generally multi-etiology

Predisposing and Risk Factors



- **Advanced age**
- **Dementia**
- **Medical comorbidity**
- **History of alcohol abuse**
- **Male sex**
- **Sensory impairments especially hearing and vision**
- **Unfamiliar settings**
- **Language barrier**
- **Dehydration**

Potential Precipitating Factors



- **Environmental Changes**
- **Pain**
- **Emotional stress or social isolation**
- **Prolonged sleep deprivation**
- **Prolonged sensory deprivation**
- **Prolonged sensory stimulation**
- **Metabolic disturbance**

Potential Precipitating Factors



- **Neurological conditions**
- **Surgery**
- **Severe medical illness or trauma**
- **Anaesthetic exposure**
- **Infections**
- **Hypoxia**
- **Fever or hypothermia**

Potential Precipitating Factors



- **Drugs and Medications**
 - Sedating hypnotics
 - Narcotics (Opiates)
 - Anticholinergic drugs
 - Cocaine and psychostimulants
 - Alcohol withdrawal
 - Sedative withdrawal

Vulnerability Liability Model




Predisposing factors and vulnerabilities

High Vulnerability

- Severe Dementia
- Post- Surgery
- Alcoholism
- Dehydration
- Elderly

• Young & Health




Precipitating Events

High Noxious Insults

- Multi-trauma
- Alcohol withdrawal
- Dehydration
- Medications
- Sleep Deprivation

• Environment Change



Vulnerability Liability Model



- High morbidity
- Increases mortality
- Difficult to diagnose even in medical settings
- In severe to extreme cases: a medical emergency
- In otherwise health individuals sudden onset delirium may be warning signs of potentially life threatening metabolic changes
- Additional insults can lead to cardiac arrest or sudden death

Managements of Delirium



- **Social restraint**
 - Familiar and low stimuli environment
 - Orienting stimuli
- **Rehydration**
- **Normalization of sleep-wake cycle**
- **Correct underlying medical conditions**
- **If restrained, pharmacological treatment should be initiated ASAP except in rare situations**
 - Benzodiazepines
 - Antipsychotic

Suggestions for First Responders



- Education, training and recognition
- Use containment whenever possible
- Medical/paramedic backup as soon as possible
- Awareness of the risks of restraints, energy weapons
- Frequent monitor of basic vitals
 - Heart rate, breathing and level of consciousness
- In “excited delirium”
 - Sedation, oxygen, IV fluid, cooling measures, low stimuli environment as soon as possible
- Available defibrillators?